arXiv:quant-ph/0101056v2 13 Feb 2002

Mesoscopic superpositions of vibronic collective states of N trapped ions

E. Solano,[1,2] R. L. de Matos Filho,[1] and N. Zagury [1]

[1] *Instituto de Física, Universidade Federal do Rio de Janeiro, Caixa Postal 68528, 21945-970 Rio de Janeiro, RJ, Brazil*

[2] *Sección Física, Departamento de Ciencias, Pontificia Universidad Católica del Perú, Apartado 1761, Lima, Peru*

(February 1, 2008)

We propose a scalable procedure to generate superpositions of motional coherent states and also entangled vibronic states in N trapped ions. Beyond their fundamental importance, these states may be of interest for quantum information processing and may be used in experimental studies on decoherence.

PACS number(s): 3.65.Ud, 42.50.Vk, 3.67.Hk

In quantum mechanics, the superposition principle, requiring that all formal superpositions of quantum states give rise to new physical states, plays a fundamental role [1]. In particular, it implies that probability densities of observable quantities, in a superposition state, usually exhibit interference effects instead of simply being added. This principle has been successfully applied to explain a large class of new phenomena in the microscopic level. As an important example one can refer to the prediction of superpositions of the K^0 meson and its antiparticle as forming new and different states, the so called K_1^0 and K_2^0 particles [2]. Superpositions of product states of composite systems also lead to the fundamental quantum nonlocality, characteristic of quantum entanglement. The quantum correlations resulting of such superpositions lead to the violation of the Bell's inequalities [3] and are essential ingredients for quantum computation [4] and quantum cryptography [5].

Despite its experimental manifestations in the microscopic level, the quantum superposition principle seems not to be applicable in the macroscopic world, as it would predict, for example, the existence of superpositions of a macroscopic object being at different places. Such situation would stay in clear contradiction to our observations and measurements of macroscopic phenomena. In fact, the great majority of states predicted by the superposition principle are not observed in our daily life. In recent years decoherence [6] is being widely accepted as the mechanism responsible for this fact and, consequently, for the emergency of classicality. It consists in the fast decay of quantum superpositions into statistical mixtures, and can be viewed as a consequence of always present interactions between a quantum system and its environment. It precludes, in general, the existence of macroscopic superpositions, except for very short time intervals and is the key to understanding the fuzzy boundary between quantum and classical behavior. As coherence disappears very fast as the system grows, it is important to be able to generate mesoscopic superposition of states and analyze how it looses its coherence. In this context, superpositions of coherent states of both the electromagnetic field in a high Q microwave cavity [8] and of the vibrational motion of a single trapped ion [9] have been generated and their behavior in the presence of the environment has been studied. Fast generation of superposition of states involving a large number of particles and excitations has become an experimental challenge in order to understand how the transition from quantum to classical occurs.

In this paper, we discuss some proposals for generating, in a fast and controllable way, vibronic superpositions of mesoscopic states involving, in principle, an arbitrary number N of ions. Relying on homogeneous resonant bichromatic excitations of the N ions with laser light, we show that is possible to generate superpositions of $N+1$ coherent states of the center of mass (CM) vibration of the N ions, equally spaced in phase space. We also show that it is possible to obtain mesoscopic entangled states involving the internal and external degrees of freedom of the N ions and odd and even coherent superpositions of the CM motion of the ions, by additionally using a dispersive bichromatic interaction. Our procedure opens the possibility of generating mesoscopic superpositions of several massive particles in a scalable and fast way.

Let us consider N two-level ions of mass m, confined to move in the z direction in a Paul trap. They are cooled down to very low temperatures [10,11] and may perform small oscillations around their equilibrium positions, $z_{j0}, j = 1, 2...N$. We denote by $Z = \sum_{j=1,N} z_j/N$, the center of mass coordinate and we set the origin at its equilibrium position. All ions are simultaneously illuminated by two classical homogeneous Raman effective pulses $\vec{E}_I = \vec{E}_{0I} e^{i(\vec{q}_1\cdot\vec{r}-\omega_I t-\varphi_I)}$ and $\vec{E}_{II} = \vec{E}_{0II}\, e^{i(\vec{q}_2\cdot\vec{r}-\omega_{II} t-\varphi_{II})}$, with angular frequencies ω_I and ω_{II} and wave vectors $\vec{q}_1 = \vec{q}_2 = \vec{q}$, parallel to the z direction. The Raman pulses frequencies will be chosen to be quasi-resonant with a long-living electronic transition between two ionic hyperfine levels $|e_j\rangle$ and $|g_j\rangle$ $(j = 1, ...N)$, with energies $\hbar\omega_0$ and 0, respectively. The Raman laser relative phases are chosen to be the same $\varphi_I = \varphi_{II} = \varphi$. The total Hamiltonian of the system may be written, in the optical rotating wave approximation (RWA), as

$$\hat{H} = \hat{H}_0 + \hat{H}_{\rm int}, \tag{1}$$

with

$$\hat{H}_0 = \hbar\omega_0 \sum_{j=1,N} |e_j\rangle\langle e_j| + \hbar\nu\hat{a}^\dagger\hat{a} + \sum_{\lambda=1,N-1} \hbar\nu_\lambda \hat{b}_\lambda^\dagger \hat{b}_\lambda\,, \tag{2}$$

and

$$\hat{H}_{\text{int}} = \hbar\Omega \sum_{j=1,N} e^{i(q\hat{z}_j-\varphi)}|e_j\rangle\langle g_j| \left(e^{-i\omega_I t} + e^{-i\omega_{II} t}\right) + \text{H.c.} \tag{3}$$

The operators $\hat{a}$ and $\hat{b}_\lambda$ ($\hat{a}^\dagger$ and $\hat{b}_\lambda^\dagger$) are the annihilation (creation) operators associated with the center of mass mode of frequency ν and with the $N-1$ other vibrational modes of frequency ν_λ, respectively. For simplicity, we have assumed that the same Rabi frequency Ω (taken as real) is associated with both lasers.

We start by taking the frequencies ω_I and ω_{II} resonant with the center of mass vibronic transition in the k-th blue and k-th red sideband as

$$\omega_I = \omega_0 + k\nu \quad \text{and} \quad \omega_{II} = \omega_0 - k\nu\,. \tag{4}$$

For small k values, we may safely assume that only the center of mass motion will be excited, given that the next eigenfrequency is $\nu_r = \sqrt{3}\nu$, corresponding to the stretch mode. The following frequencies ($\geq \sqrt{29/5}\nu$)depend on the number of ions and have being calculated in Ref. [12]. Following the usual treatment for one single ion interacting with a laser field [13], we make the RWA with respect to the CM vibrational frequency and select the terms that oscillate with minimum frequency. In the Lamb-Dicke limit the interaction Hamiltonian may be written, in the interaction picture, as

$$H_{\text{int}} = \frac{2\hbar\Omega\eta^k}{k!}\hat{J}_{\text{T}}(\hat{a}^k + \hat{a}^{\dagger k}) \tag{5}$$

where $\eta = q\sqrt{\hbar/2Nm\nu}$ is the Lamb-Dicke parameter associated to the center of mass motion of the N ions, $\hat{J}_{\text{T}}$ is an angular momentum-like operator [14] defined as

$$\hat{J}_{\text{T}} = \frac{i^k e^{-i\varphi}}{2} \sum_{j=1,N} \hat{\sigma}_{j+} + \text{H.c.} \tag{6}$$

Here $\widehat{\sigma}_{+j} = |\uparrow_j\rangle\langle\downarrow_j| = e^{iqz_{j0}}|e_j\rangle\langle g_j|$ is a flip operator associated with the electronic transition $|g_j\rangle \rightarrow |e_j\rangle$ in the ion j and $\widehat{\sigma}_{j-} = \widehat{\sigma}_{j+}^\dagger$. Without loss of generality we may set the phase $\varphi = k\pi/2$, so that $\hat{J}_{\text{T}} = \hat{J}_x$ in the usual angular momentum operator convention for phases. Similarly, by choosing a phase $\varphi = (k+1)\pi/2$, $\hat{J}_{\text{T}} = \hat{J}_y$. We also may define the z component of the angular momentum by $\hat{J}_z = \frac{1}{2}\sum_j(|e_j\rangle\langle e_j| - |g_j\rangle\langle g_j|)$.

From Eq. (5), it is easy to show that the time evolution operator, in the interaction picture, at time t, is a sum of products of unitary operators on the motional states and projection operators on the ion internal states

$$\hat{U}_k(t) = \sum_{j,m} \hat{D}_k(m\alpha_k(t))|j,m\rangle_{x\,x}\langle j,m| \tag{7}$$

where

$$\hat{D}_k(\alpha_k) = e^{\alpha_k \hat{a}^{k\dagger} - \alpha_k * \hat{a}^k}\,, \tag{8}$$

with $\alpha_k(t) = 2i\Omega t\eta^k/k!$. Also, $|j,m\rangle_x$ are the simultaneous eigenvectors of the operators $\hat{J}_x$ and $\hat{J}^2 \equiv \hat{J}_x^2 + \hat{J}_y^2 + \hat{J}_z^2$, associated with the eigenvalues $m = -j, -(j-1), ...j$ and $j(j+1)$, respectively. j varies from 0 (1/2) to $N/2$ by steps of 1, if N is even (odd).

The action of the time evolution operator $\hat{U}_k$ of Eq. (7) corresponds to unitary operations $\hat{D}_k(m\alpha_k)$ on the motional degrees of freedom conditioned to the value m of the x component of the "angular momentum" electronic state. From now on we set $k = 1$ that is, the excitation occurs in the first red and blue sidebands. In this case $\hat{D}_1(\alpha)$ is the displacement operator which generates, when acting on the ground state, coherent states of the vibrational motion of the center of mass

$$|\alpha\rangle_{\text{coh}} = e^{(\alpha\hat{a}^\dagger - \alpha * \hat{a})}|0\rangle = e^{-|\alpha|^2/2}\sum \frac{\alpha^n}{\sqrt{n!}}|n\rangle\,. \tag{9}$$

If initially the ions are in the vibronic ground state, $|ggg...\rangle \otimes |0\rangle \equiv |N/2, -N/2\rangle_z \otimes |0\rangle$, their state, after an interaction time t with the laser fields, will be given by

$$\sum_{m,m'} d^{N/2}_{m',m}(\pi/2) d^{N/2}_{m,-N/2}(-\pi/2)|N/2, m'\rangle_z \otimes |m\alpha\rangle_{\text{coh}}, \tag{10}$$

where $\alpha = 2i\eta\Omega t$. Here, $d^j_{m',m}(\theta) = {}_z\langle j,m'|e^{-i\theta J_y}|j,m\rangle_z$ are the matrix elements of the rotation operator along the y axis in the $\{|j,m\rangle_z\}$ basis of the eigenstates of $\hat{J}_z$ and $\hat{J}^2$ [16]. This allows us to prepare in a very simple way and with a single Raman pulse, a mesoscopic superposition of vibronic quantum states in N trapped ions. If we now measure the electronic state of the ions and find the totally excited state $|eee...\rangle$, we know that the motional state is given, up to a normalization factor, by

$$\sum_{m=-N/2,N/2} \frac{(-1)^{(N/2-m)}}{(N/2-m)!(N/2+m)!}|m\alpha\rangle_{\text{coh}}. \tag{11}$$

If N is even, this state is a superposition of the vacuum and a series of coherent states of amplitudes $m\alpha$, the probability for measuring them decreasing with $|m|$. If N is odd, the vacuum is not present in the superposition. When we have only one ion Eq. (11) represents a single odd coherent state. Note that the procedure above relies on a resonant excitation of the ions and therefore can be accomplished with very *short* interaction times. The measurement of the electronic state can be done by monitoring the fluorescence of a cyclic transition of the ions [17], where a dark event detects the totally excited state. In Fig. 1, we show the Wigner function for the state given by Eq. (11), when $|\alpha| = 3$ and $N = 3$. Note that the vacuum is not present, as N is odd, and that interference manifests strongly, due to the multiple superposition of coherent states.



FIG. 1. Wigner distribution for the state given in Eq. (11) when $N = 3$ and $|\alpha| = 3$.

If the frequencies of the lasers in Eq. (3) are slightly off resonance and such that $\omega_I = \omega_0 + \nu + \delta$ and $\omega_{II} = \omega_0 - \nu - \delta$, the time evolution operator may be written, for small values of η, approximately as $e^{-i\lambda t \hat{J}_y^2}$, where $\lambda = 8(\Omega\eta)^2/\delta$ and we have chosen $\varphi = 0$. This dispersive bichromatic interaction does not change the motional state and has been studied in a series of recent papers [18–21]. It has been implemented successfully in the laboratory for generating GHZ states in two and four ions [11].

We now show that, by using both the dispersive and the resonant bichromatic interaction, it is possible to prepare even and odd coherent states of large amplitudes and involving a large number of ions. It will be convenient to consider first the case where we have an odd number of ions N. If we start with the ground state $|ggg...\rangle \otimes |0\rangle$ and apply the dispersive bichromatic interaction during a time $t = \pi/(2\lambda)$, we get, using the properties of the rotation matrices [16],

$$e^{-i\hat{J}_y^2 \pi/2}|j,-j\rangle_z \otimes |0\rangle = \frac{1}{\sqrt{2}}(|j,j\rangle_x - |j,-j\rangle_x) \otimes |0\rangle \,, \tag{12}$$

where $j = N/2$.

Now we apply a second pulse using the resonant lasers of frequencies $\omega_0 \pm \nu$, with a relative phase of $\pi/2$ with respect to the previous pulse, during a time τ. This corresponds to apply the operator $U_1(\tau)$ on the state given in Eq. 12 leading to the state

$$\begin{aligned} &\tfrac{1}{\sqrt{2}}(|N\alpha/2\rangle_{\rm coh} \otimes |N/2, N/2\rangle_x \\ &- |-N\alpha/2\rangle_{\rm coh} \otimes ||N/2, -N/2\rangle_x \,. \end{aligned} \tag{13}$$

where $\alpha = 2i\Omega\eta\tau$. This state is a superposition of two vibronic states involving, in principle, a large number N of ions. The two internal states $|N/2, \pm N/2\rangle_x$ are strongly correlated to the two vibrational states $|\pm N\alpha/2\rangle_{\rm coh}$, whose average amplitude of oscillation, $|N\alpha/2|$, is proportional both to $\sqrt{N}$ and $\Omega\tau$. This state is an example of a strongly entangled mesoscopic state that is scalable, and its experimental realization may become a useful tool in analyzing the dependence of decoherence on the number of degrees of freedom of this system.

If, after the preparation of the state given in Eq. (13), we measure the state $|eee...\rangle$, we obtain

$$\frac{|N\alpha/2\rangle_{\rm coh} + |-N\alpha/2\rangle_{\rm coh}}{\sqrt{2 + 2e^{-|N\alpha|^2/2}}} \,. \tag{14}$$

The state given in Eq. (14) is an even coherent state of the center of mass motion of an odd large number of ions. An odd coherent state is obtained if one measures the state $|ggg...\rangle$ instead of $|eee...\rangle$. Thus, we have a procedure that generates a superposition of two states of motion, vibrating out of phase, where a large number of particles may be involved. In principle, the number of ions in this state is limited mostly by decoherence and by the capacity of producing lasers with high homogeneity. For large N, the probability to produce these state superpositions by this procedure decreases as $1/2^N$. However, an efficient method to obtain even or odd coherent states, each one with probability 1/2, even for large N, will be described below.

As before, we start with the ground state and apply in succession the dispersive bichromatic interaction as in Eq. (12) and the resonant bichromatic interaction to obtain the state given in Eq. (13). We then apply again the same dispersive bichromatic interaction during the same interval of time. The resulting state is

$$\begin{aligned} &\frac{1}{2}\{|eee..\rangle \otimes (|N\alpha/2\rangle - |-N\alpha/2\rangle) \\ &-|ggg...\rangle \otimes (|N\alpha/2\rangle + |-N\alpha/2\rangle)\} \end{aligned} \tag{15}$$

Measuring the dark fluorescence of either the $N-$ electronic excited state or the $N-$ electronic ground state, we get the associated odd or even coherent state with probability 1/2. This procedure shows how to generate efficiently mesoscopic superposition states for a large number of ions.

Similar results may be obtained for N even, if an additional resonant carrier $\pi/2$ pulse, is applied at the same time and with a relative phase $\varphi = \pi/2$ with respect to the dispersive pulse.

In conclusion, we have presented a procedure to generate several kinds of mesoscopic superpositions of states involving N ions. Superpositions of coherent states evenly spaced on a line may be generated rapidly, through resonant interactions. Even or odd coherent state may be obtained if we use also dispersive bichromatic interactions. This is, to the best of our knowledge, the first proposal to generate, in a scalable way, mesoscopic superpositions of collective motional states of N trapped ions. Note that, as long as the field is spatially homogeneous over the trapped ions, the only parameter that depends on N, in our model, is the time we applied the laser pulses, which varies with $\sqrt{N}(N)$ during the resonant (dispersive) interaction. We also expect that the cooling of the vibrational modes for N ions will be achieved with the same effectiveness as in the case of four ions [11]. For this reason the constraints, in our proposal, on parameters like laser pulse timing, laser frequencies, as well as cooling to the ground state should remain almost the same as in the case of a single trapped ion. Then, the number of ions involved will be limited mostly by the scale of time in which decoherence occurs.

We believe that the proposals presented in this letter are ready to be implemented in the laboratory. They should help to build larger mesoscopic quantum superpositions in trapped ions, to study experimentally, at large scale, decoherence processes and for applications in quantum information processing.

This work was partially supported by the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), the Fundação de Amparo a Pesquisa do Estado do Rio de Janeiro (FAPERJ), the Programa de Apoio a Núcleos de Excelência (PRONEX) and Fundação José Bonifácio (FUJB).

[1] P. A. M. Dirac, *The Principles of Quantum Mechanics*, Claredon Press, Oxford (1947).

[2] M. Gell-Mann and A. Pais, Phys. Rev. **97**, 1387 (1955); T. D. Lee and C. N. Yang, Phys. Rev. **104**, 254 (1956).

[3] J. S. Bell, Physics **1**, 195 (1964).

[4] D. Deutsch, Proc. R. Soc. London A **400**, 97 (1985); A. Berthiaume and G. Brassard, in Proceedings of the Seventh Annual IEEE Conference on Structure in Complexity Theory, Boston, June 1992 (IEEE, New York, 1989), p. 132.

[5] S. Wiesner, Sigact News **15**, 78 (1983); A. K. Ekert, Phys. Rev. Lett. **67**, 661 (1991); C. H. Benett, G. Brassard, and N. D. Mermin, *ibid* **68**, 557 (1992); C. H. Benett, *ibid* **68**, 3121 (1992); A. K. Ekert, J. G. Rarity, P. R. Tapster, and G. M. Palma, *ibid* **69**, 1293 (1992).

[6] W. H. Zurek Physics Today **44**, 36 (1991); Roland Omnès in *The Interpretaion of Quantum Mechanics*, Princeton University Press, Princeton, 1994; D. Giulini, E. Joos, C. Kiefer, J. Kupsch, I.-O. Stamatescu, and H. D. Zeh, *Decoherence and the Appearance of a Classical Word in Quantum Theory*, Springer-Verlag, Berlin, Heidelberg (1996); J. P. Paz and W. H. Zurek *Environment-induced decoherence and the transition from quantum to classical*, LXXII Les Houches Summer School, 1999.

[7] E. Schrödinger, Naturwissenschaften **23**, 807(1935).

[8] M. Brune, E. Hagley, J. Dreyer, X. Maître, A. Maali, C. Wunderlich, J. M. Raimond, and S. Haroche, Phys. Rev. Lett. **77**, 4887(1996).

[9] C. Monroe, D. M. Meekhof, B. E. King, and D. J. Wineland, Science **272**, 1131 (1996); C. J. Myatt, B. E. King, Q. A. Turchette, C. A. Sackett, D. Kielpinski, W. M. Itano, C. Monroe, and D. J. Wineland, Nature,**403**,269 (2000); Q. A. Turchette, C. J. Myatt, B. E. King, C. A. Sackett, D. Kielpinski, W. M. Itano, C. Monroe, and D. J. Wineland, Phys. Rev. A **62**, 053807 (2000).

[10] W. Neuhauser, M. Hohenstatt, P. Toschek, and H. Dehmelt, Phys. Rev. Lett. 41, 233 (1978); B. E. King, C. S. Wood, C. J. Myatt, Q. A. Turchette, D. Leibfried, W. M. Itano, C. Monroe, and D. J. Wineland, Phys. Rev. Lett. **81**, 1525 (1998); C. F. Roos, D. Leibfried, A. Mundt, F. SchmidtKaler, J. Eschner, and R. Blatt, Phys. Rev. Lett. 85, 547 (2000).

[11] C. A. Sackett, D. Kielpinski, B. E. King, C. Langer, V. Meyer, C. J. Myatt, M. Rowe, Q. A. Turchette, W. M. Itano, D. J. Wineland and C. Monroe, Nature **404**, 256 (2000).

[12] D. F. V. James, Appl. Phys. B **66**, 181 (1998).

[13] W. Vogel and R.L. de Matos Filho, Phys. Rev. A**52**, 4214 (1995).

[14] see for example H. M. Nussenzveig, in Introduction to Quantum Optics, Gordon and Breach, New York, 1973, pgs. 174-176

[15] H. P. Yuen, Phys. Rev. A**13**, 2226 (1971).

[16] see for example K. Gottfried, in Quantum Mechanics, W. A. Benjamin, New York, 1966, pgs. 264-288.

[17] W. Nagourney and J. Sandberg and H. Dehmelt, Phys. Rev. Lett., **56**, 2797 (1986).

[18] A. Sørensen and K. Mølmer, Phys. Rev. Lett. **82**, 1971 (1999).

[19] K. Mølmer and A. Sørensen, Phys. Rev. Lett.**82**,1835 (1999).

[20] A. Sørensen and K. Mølmer, Phys. Rev. A, **62**, 022311 (2000).

[21] E. Solano, R. L. de Matos Filho, and N. Zagury, Phys. Rev. A, **59**, R2539 (1999); **61**, 029903(E) (2000).

[22] Q. A. Turchette, C. J. Myatt, B. E. King, C. A. Sackett, D. Kielpinski, W. M. Itano, C. Monroe, and D. J. Wineland, Phys. Rev. A, **62**, 053807 (2000).